



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Addendum

**Citation for published version:**

Howard, DM, Adams, MJ, Shirali, M, Clarke, T-K, Marioni, RE, Davies, G, Coleman, JRI, Alloza, C, Shen, X, Barbu, MC, Wigmore, EM, Gibson, J, Hagenaars, SP, Lewis, CM, Ward, J, Smith, DJ, Sullivan, PF, Haley, CS, Breen, G, Deary, IJ & McIntosh, AM 2018, 'Addendum: Genome-wide association study of depression phenotypes in UK Biobank identifies variants in excitatory synaptic pathways', *Nature Communications*, vol. 9, no. 1, pp. 3578. <https://doi.org/10.1038/s41467-018-05310-5>

**Digital Object Identifier (DOI):**

[10.1038/s41467-018-05310-5](https://doi.org/10.1038/s41467-018-05310-5)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Nature Communications

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



DOI: 10.1038/s41467-018-05310-5

OPEN

# Addendum: Genome-wide association study of depression phenotypes in UK Biobank identifies variants in excitatory synaptic pathways

David M. Howard<sup>1</sup>, Mark J. Adams<sup>1</sup>, Masoud Shirali<sup>1</sup>, Toni-Kim Clarke<sup>1</sup>, Riccardo E. Marioni<sup>2</sup>, Gail Davies<sup>2,3</sup>, Jonathan R.I. Coleman<sup>4,5</sup>, Clara Alloza<sup>1</sup>, Xueyi Shen<sup>1</sup>, Miruna C. Barbu<sup>1</sup>, Eleanor M. Wigmore<sup>1</sup>, Jude Gibson<sup>1</sup>, 23andMe Research Team<sup>#</sup>, Saskia P. Hagenaars<sup>4,5</sup>, Cathryn M. Lewis<sup>4,5</sup>, Joey Ward<sup>6</sup>, Daniel J. Smith<sup>6</sup>, Patrick F. Sullivan<sup>7,8,9</sup>, Chris S. Haley<sup>10</sup>, Gerome Breen<sup>4,5</sup>, Ian J. Deary<sup>2,3</sup> & Andrew M. McIntosh<sup>1,3</sup>

Addendum to: *Nature Communications*; <https://doi.org/10.1038/s41467-018-03819-3>; published online 16 Apr 2018

In this article, a meta-analysis of the significant variants in the UK Biobank sample with a previously published study by 23andMe was included. Due to the individual study results being reported on different scales, the meta-analytic point estimates were not reliable. We have re-analysed the 17 significant variants in the UK Biobank using a comparable approach to 23andMe in Plink. We present an updated version of Table 1 associated with this Addendum that includes the newly calculated comparable effect size estimates for UK Biobank, the updated meta-analysis results and we have extended the table legend to provide information on the updated columns. As a consequence of the new analysis, the original statement in the first paragraph under the subheading “Genome-wide association study of depression” reading “All 17 variants remained significant ( $P < 5 \times 10^{-8}$ ) in the meta-analysis” should be amended to “10 variants remained significant ( $P < 5 \times 10^{-8}$ ) in the meta-analysis”. There was no change to the direction of allelic effect in either cohort.

The fourth sentence in Methods section under the subheading “Replication cohort and meta-analysis” on page 8 was incomplete and should read “Additionally, we used Metal<sup>36</sup> to conduct an inverse variance-weighted meta-analysis, using LD score regression intercepts<sup>13</sup> for genomic inflation control.”, adding “using LD score regression intercepts<sup>13</sup> for genomic inflation control” at the end of the sentence.

We have also included transformed effect sizes and standard errors in our summary statistics deposited on the Edinburgh DataShare website and have updated the doi to <https://doi.org/10.7488/ds/2350>.

In a second related issue, we used the new BGENIE software package to generate the results with which to conduct gene, region and gene-set analyses to generate data presented in the “Gene and region-based analysis” and “Gene-set pathway analysis” sections, Table 2, Supplementary Table 6, Supplementary Data 6–12, and Supplementary Figures 7–9. BGENIE reported minor allele frequencies (MAF) across the whole of UK Biobank ( $n = 487,409$ ) rather than based on those individuals that were included in each of the association analyses (broad depression  $n = 322,580$ ; probable major depressive disorder  $n = 174,519$ ; International Classification of Diseases-coded major depressive disorder  $n = 217,584$ ). Therefore the results reported in the

<sup>1</sup> Division of Psychiatry, University of Edinburgh, EH10 5HF Edinburgh, UK. <sup>2</sup> Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, EH8 9JZ Edinburgh, UK. <sup>3</sup> Department of Psychology, University of Edinburgh, EH8 9JZ Edinburgh, UK. <sup>4</sup> Social Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King’s College London, SE5 8AF London, UK. <sup>5</sup> NIHR Biomedical Research Centre for Mental Health, South London and Maudsley NHS Trust, SE5 8AF London, UK. <sup>6</sup> Institute of Health and Wellbeing, University of Glasgow, G12 8RZ Glasgow, UK. <sup>7</sup> Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, 171 77 Stockholm, Sweden. <sup>8</sup> Department of Genetics, University of North Carolina, 27599 Chapel Hill, NC, USA. <sup>9</sup> Department of Psychiatry, University of North Carolina, 27599 Chapel Hill, NC, USA. <sup>10</sup> Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, EH4 2XU Edinburgh, UK. \*A full list of consortium members appears at the end of the paper. Correspondence and requests for materials should be addressed to D.M.H. (email: [D.Howard@ed.ac.uk](mailto:D.Howard@ed.ac.uk))

“Gene and region-based analysis” and “Gene-set pathway analysis” were based on the MAF across the whole of UK Biobank as opposed to what we described in the methods section. To correct these errors in our initial analysis, we have now re-analysed the data using the MAF based on only those individuals included in the respective association analysis. This has resulted in 73 rather than 78 significant genes for broad depression, three rather than two genes for probable MDD, and zero genes rather than one gene for ICD-coded MDD which was originally described in the “Gene and region-based analysis” section in the results. The gene-based results described on page 3 under the subheading “Gene and region-based analyses” should read “We used the MAGMA<sup>18</sup> package to identify genes with a significant effect ( $P < 2.77 \times 10^{-6}$ ) on each phenotype. There were 73 genes significantly associated with broad depression (Supplementary Data 6, associated with this Addendum), and three genes that were associated with probable MDD (Supplementary Data 7, associated with this Addendum)”. Updated versions of Supplementary Note 1, Supplementary Figures 7, 8, and 9 in the Supplementary Information file, and Supplementary Data 6 and 7 are included here to reflect these changes in the data. The original Supplementary Data 8 should be disregarded as with the new analysis no genome-wide significant SNPs for ICD-coded MDD are detected. In the region-based analysis described on page 3 number of regions assessed increased from 8308 to 8345 requiring a slight change in the significance threshold from  $6.02 \times 10^{-8}$  to  $5.99 \times 10^{-6}$  so that the sentence in the second paragraph under the subheading “Gene and region-based analyses” now reads “We also used MAGMA to identify genomic regions, defined by recombination hotspots, with a statistically significant effect ( $P < 5.99 \times 10^{-6}$ ) on each phenotype”. The number of significant regions remained the same and updated estimates of effect sizes of regions may be found in the updated Supplementary Data 8, 10, and 11 associated with this Addendum. The significant gene-sets described under the subheading “Gene-set pathway analysis” remained the same with slight differences in number of genes in each pathway, effect sizes and  $p$ -values. The first sentence in this section now reads: “We conducted gene-set enrichment analysis<sup>19,20</sup> and identified five significant pathways for broad depression after applying correction for multiple testing: GO\_EXCITATORY\_SYNAPSE (beta =  $0.342 \pm 0.069$ ,  $P_{corrected}$  = 0.003), GO\_MECHANOSENSORY\_BEHAVIOR (beta =  $1.270 \pm 0.294$ ,  $P_{corrected}$  = 0.047), GO\_POSTSYNAPSE (beta =  $0.248 \pm 0.050$ ,  $P_{corrected}$  = 0.003), GO\_NEURON\_SPINE (beta =  $0.390 \pm 0.089$ ,  $P_{corrected}$  = 0.019) and GO\_DENDRITE (beta =  $0.200 \pm 0.044$ ,  $P_{corrected}$  = 0.021) (Table 2)”. Amended versions of Table 2, Supplementary Data 12 and Supplementary Table 6 in the Supplementary Information file are included in this Addendum.

Table 1 Independent variants with a genome-wide significant ( $P < 5 \times 10^{-8}$ ) association with broad depression, probable major depressive disorder (MDD), or International Classification of Diseases (ICD)-coded MDD in the UK Biobank															
Phenotype	Chr	Marker name	Position	UK Biobank					23andMe			Meta-analysis			
				A1/A2	Freq	Info	log OR (St Err)	P-value	Gene $\pm$ 10 kb	log OR (St Err)	P-value	log OR (St Err)	P-value	Direction	
Broad depression	1	rs10127497	67050144	T/A	0.138	1.00	0.044 (0.0077)	$1.26 \times 10^{-8}$	SGIP1	0.0098 (0.0086)	0.257	0.029 (0.0058)	$5.70 \times 10^{-7}$	++	
	1	rs6699744	72825144	T/A	0.612	1.00	0.040 (0.0055)	$1.64 \times 10^{-13}$	—	0.0328 (0.0064)	$2.68 \times 10^{-7}$	0.0372 (0.0042)	$9.76 \times 10^{-19}$	++	
	1	rs6424532	73664022	A/G	0.486	1.00	0.029 (0.0053)	$3.74 \times 10^{-8}$	—	0.0233 (0.006)	$1.14 \times 10^{-4}$	0.0267 (0.004)	$4.10 \times 10^{-11}$	++	
	1	rs7548151	177026983	A/G	0.084	1.00	0.056 (0.0095)	$3.77 \times 10^{-9}$	ASTN1	0.006 (0.0104)	0.560	0.0339 (0.0071)	$1.97 \times 10^{-6}$	++	
	5	rs40465	103981726	G/T	0.332	1.00	0.035 (0.0056)	$4.37 \times 10^{-10}$	RP11-6N13.1	0.0193 (0.0064)	$2.63 \times 10^{-3}$	0.0285 (0.0043)	$3.53 \times 10^{-11}$	++	
	6	rs3132685	29945949	A/G	0.130	1.00	−0.059 (0.0081)	$2.48 \times 10^{-13}$	—	−0.0249 (0.0099)	0.0115	−0.0459 (0.0063)	$4.87 \times 10^{-13}$	--	
	6	rs112348907	73587953	G/A	0.296	1.00	0.033 (0.0058)	$1.44 \times 10^{-8}$	—	−0.0004 (0.0067)	0.950	0.019 (0.0045)	$2.06 \times 10^{-5}$	++	
	7	rs3807865	12250402	A/G	0.412	1.00	0.037 (0.0054)	$6.81 \times 10^{-12}$	TMEM106B	0.019 (0.0061)	0.002	0.0293 (0.0041)	$8.79 \times 10^{-13}$	++	
	7	rs2402273	117600424	C/T	0.409	1.00	0.032 (0.0054)	$1.88 \times 10^{-9}$	—	0.0093 (0.0061)	0.130	0.0226 (0.0041)	$3.97 \times 10^{-8}$	++	
	9	rs263575	17033840	A/G	0.460	1.00	−0.030 (0.0053)	$2.36 \times 10^{-8}$	—	−0.0157 (0.0061)	$9.45 \times 10^{-3}$	−0.0238 (0.0041)	$4.51 \times 10^{-9}$	--	
	10	rs1021363	106610839	G/A	0.642	1.00	−0.032 (0.0055)	$1.02 \times 10^{-8}$	SORCS3	−0.031 (0.0063)	$9.34 \times 10^{-7}$	−0.0314 (0.0042)	$1.04 \times 10^{-13}$	--	
	11	rs10501696	88748162	G/A	0.499	0.99	−0.036 (0.0054)	$6.42 \times 10^{-11}$	GRM5	−0.0251 (0.0066)	$1.49 \times 10^{-4}$	−0.0315 (0.0043)	$1.55 \times 10^{-13}$	--	
	13	rs9530139	31847324	T/C	0.195	1.00	−0.040 (0.0067)	$2.59 \times 10^{-9}$	B3GLCT	−0.0075 (0.0078)	0.338	−0.0265 (0.0052)	$2.77 \times 10^{-7}$	--	
	15	rs28541419	88945878	G/C	0.231	1.00	−0.035 (0.0064)	$2.82 \times 10^{-8}$	—	−0.0029 (0.0073)	0.688	−0.0218 (0.0049)	$7.75 \times 10^{-6}$	--	
Probable MDD	2	rs10929355	15398964	G/T	0.456	1.00	−0.053 (0.0090)	$5.89 \times 10^{-9}$	NBAS	−0.0078 (0.0061)	0.199	−0.0221 (0.0051)	$1.62 \times 10^{-5}$	--	
ICD-coded MDD	7	rs5011432	12268668	C/A	0.412	1.00	0.051 (0.0091)	$2.10 \times 10^{-8}$	TMEM106B	0.022 (0.0061)	$3.15 \times 10^{-4}$	0.0313 (0.0052)	$1.47 \times 10^{-9}$	++	
MDD	7	rs1554505	1983929	A/G	0.752	1.00	0.114 (0.0191)	$2.73 \times 10^{-9}$	MAD1L1	0.017 (0.007)	0.015	0.0291 (0.0068)	$1.63 \times 10^{-5}$	++	

The UK Biobank results are from logistic regression analyses in PLINK of variants identified as genome-wide significant using BGENIE. Variants were examined within the 23andMe association analysis of depression<sup>4</sup> to obtain their reported *P*-values and determine whether their effect was in the same direction as UK Biobank. The allele frequency (Freq) is for the A1 allele within UK Biobank, with the effect size (Log OR) and standard error (St. err.) reported for the A1 allele within UK Biobank, 23andMe and the meta-analysis. The chromosome (Chr) and basepair position is given for the GRCh37 assembly. Imputation accuracy (Info) score of UK Biobank was calculated based on the sample analysed

Table 2 Pathways with a significant effect ( $P_{corrected} < 0.05$ ) on broad depression following multiple testing correction identified through gene-set enrichment analysis					
Phenotype	Pathway	Number of genes	Beta (St. Err.)	P-value	$P_{Corrected}$
Broad depression	GO_EXCITATORY_SYNAPSE	184	0.342 (0.069)	$3.45 \times 10^{-7}$	0.003
	GO_POSTSYNAPSE	354	0.248 (0.050)	$3.88 \times 10^{-7}$	0.003
	GO_NEURON_SPINE	115	0.390 (0.089)	$2.72 \times 10^{-6}$	0.019
	GO_DENDRITE	425	0.200 (0.044)	$3.08 \times 10^{-6}$	0.021
	GO_MECHANOSENSORY_BEHAVIOR	12	1.270 (0.294)	$8.04 \times 10^{-6}$	0.047

The described errors have not been fixed in the original article. We reaffirm that the changes do not change the main conclusions of the manuscript.

Published online: 30 August 2018

## References

4. Hyde, C. L. et al. Identification of 15 genetic loci associated with risk of major depression in individuals of European descent. *Nat. Genet.* **48**, 1031–1036 (2016).
13. Bulik-Sullivan, B. K. et al. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).
17. Middeldorp, C. M. et al. A genome-wide association meta-analysis of attention-deficit/hyperactivity disorder symptoms in population-based paediatric cohorts. *J. Am. Acad. Child. Adolesc. Psychiatry* **55**, 896–905 (2016).
18. de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS. Comput. Biol.* **11**, e1004219 (2015).
19. The Gene Ontology Consortium. Expansion of the gene ontology knowledgebase and resources. *Nucleic Acids Res.* **45**, D331–D338 (2017).
20. Subramanian, A. et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl Acad. Sci. USA* **102**, 15545–15550 (2005).
36. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. **26**, 2190–2191 (2010).

## Additional information

**Supplementary Information** accompanies this paper at <https://doi.org/10.1038/s41467-018-05310-5>.

## 23andMe Research Team

Michelle Agee<sup>11</sup>, Babak Alipanahi<sup>11</sup>, Adam Auton<sup>11</sup>, Robert K. Bell<sup>11</sup>, Katarzyna Bryc<sup>11</sup>, Sarah L. Elson<sup>11</sup>, Pierre Fontanillas<sup>11</sup>, Nicholas A. Furlotte<sup>11</sup>, David A. Hinds<sup>11</sup>, Karen E. Huber<sup>11</sup>, Aaron Kleinman<sup>11</sup>, Nadia K. Litterman<sup>11</sup>, Jennifer C. McCreight<sup>11</sup>, Matthew H. McIntyre<sup>11</sup>, Joanna L. Mountain<sup>11</sup>, Elizabeth S. Noblin<sup>11</sup>, Carrie A.M. Northover<sup>11</sup>, Steven J. Pitts<sup>11</sup>, J. Fah Sathirapongsasuti<sup>11</sup>, Olga V. Sazonova<sup>11</sup>, Janie F. Shelton<sup>11</sup>, Suyash Shringarpure<sup>11</sup>, Chao Tian<sup>11</sup>, Joyce Y. Tung<sup>11</sup>, Vladimir Vacic<sup>11</sup> & Catherine H. Wilson<sup>11</sup>

<sup>11</sup>23andMe, Inc., Mountain View, CA 94041, USA



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018